arXiv:math/0210048v3 [math.AG] 27 Apr 2005

FAMILIES OF D-MINIMAL MODELS AND APPLICATIONS TO 3-FOLD DIVISORIAL CONTRACTIONS

NIKOLAOS TZIOLAS

Abstract

Let X/T be a one parameter family of canonical 3-folds and let D be a Weil divisor on it flat over T. We study the problem of when the D_t-minimal models of X_t form a family and we obtain conditions for this to happen. As an application of this we classify terminal divisorial contractions $E \subset Y \longrightarrow \Gamma \subset X$ contracting an irreducible surface E onto the smooth curve Γ, in the case when the general section of X through Γ is a D_5 DuVal singularity.

0. *Introduction*

Let X be a 3-fold with canonical singularities and let D be a Weil divisor on it. It is known [**7**] that the sheaf of graded $\mathcal{O}_X$ algebras $R(X,D) = \oplus_{m\geq 0}\mathcal{O}_X(mD)$ is finitely generated. Moreover this is equivalent to the existence of a birational morphism $Z \xrightarrow{g} X$ which is an isomorphism in codimension 1 and such that $g_*^{-1}D$ is $\mathbb{Q}$-Cartier and g-ample. Such a morphism is unique and we call it the D-minimal model of X. The process of making a Weil divisor $\mathbb{Q}$-Cartier is very important in the three dimensional minimal model program and it is of interest to get higher dimensional analogues.

In the first part of this paper we investigate whether this can be done in families of canonical 3-folds. More precisely, we study the following problem. Let X/T be a one parameter family of canonical 3-folds and let D be a Weil divisor in X that is flat over T. Let $Z_t \xrightarrow{g_t} X_t$ be the D_t-minimal model of X_t, which exists as mentioned earlier. We want to know if they form a family, i.e., if there is a morphism $Z/T \xrightarrow{g} X/T$ whose fibers over T are the D_t-minimal models of X_t.

A special case of this problem is the following. Let $Y/T \xrightarrow{f} X/T$ be a projective morphism between families of canonical 3-folds and assume that $Y_t \xrightarrow{f_t} X_t$ is a flopping contraction for all t. Do the flops of f_t form a family? More precisely, is there a projective morphism $Y'/T \xrightarrow{f'} X/T$ such that $Y'_t \xrightarrow{f'_t} X_t$ is the flop of f_t, for all t? This problem is a special case of the previous one since Y' is the D-minimal model of X, where $D = f_*(-A)$, and A is an f-ample divisor.

This last problem is directly related to the problem of whether the minimal model program runs in families of canonical 3-folds. It does work in families of terminal 3-folds [**14**] but for families of canonical 3-folds it is still unknown if it works.

Example 2.16 shows that in general, D-minimal models do not form families. Therefore it is important to find conditions that need to be satisfied for the D-minimal models to form families. Theorems 2.4 and 2.7 show that the D_t minimal models form a family if $e(X_t; D_t)$, the number of crepant divisors of X_t with center

---

2000 *Mathematics Subject Classification* 14E30 (primary), 14E35 (secondary)..

in suppD_t, is independent of t. Corollary 2.5 gives conditions for the flops to form families. However these conditions are probably too strong and the case of families of flops deserves a much more careful study.

As an application of the one parameter theory that was developed in the first part of this paper we study and classify certain 3-fold divisorial contractions. Divisorial contractions is one of the two fundamental birational maps that appear in the three dimensional minimal model program. The other ones are the flips. Flips are known to exist and they have been classified [**14**]. The structure of divisorial contractions is still not well understood and a detailed knowledge of them will have many applications and will contribute to the understanding of the birational structure of Mori fiber spaces through the Sarkisov program.

Let $E \subset Y \xrightarrow{f} \Gamma \subset X$ be a 3-fold divisorial contraction such that X and Y have terminal singularities. In the case when Γ is a point, there are classification results due to Corti [**2**] and Kawakita [**5**], [**6**]. The case when Γ is a curve has been studied by Kawamata [**8**] and myself [**20**]. Suppose that Γ is a curve. If a terminal divisorial contraction f exists, then the general hyperplane section S of X through Γ is DuVal [**14**]. In [**20**] it is shown that given a smooth curve Γ in a terminal 3-fold X, there is always a divisorial contraction f with Y canonical. In this paper we classify divisorial contractions in the case that the general hyperplane section S of X through Γ is a D_5 DuVal singularity and Γ is a smooth curve. I believe that the general D_{2n+1} case can be treated in principle with same method that treats the D_5 case but it is much harder and the calculations involved are more complicated. The case when the general hyperplane section S of X through Γ is a D_{2n} DuVal singularity was studied in [**20**] and all such contractions were classified. Moreover, it was shown that the D_{2n} cases are very different from the D_{2n+1} cases.

Let $0 \in X$ be an index one 3-fold terminal singularity and Γ a smooth curve in X passing through the singular point. A divisorial contraction $E \subset Y \xrightarrow{f} \Gamma \subset X$ contracting an irreducible divisor onto Γ can be constructed as follows [**20**]. Let $g\colon W \longrightarrow X$ be the blow up of X along Γ. Then there are two g-exceptional divisors E and $F \cong \mathbb{P}^2$. E is a ruled surface over Γ and F is over the singularity of X. Let $h\colon Z \longrightarrow W$ be the E-minimal model of W which exists [**7**]. Then after running a suitable minimal model program we arrive to a Z' and the birational transform of F in Z' can be contracted to get the required contraction $Y \xrightarrow{f} X$.

In order to understand when a terminal contraction exists, it is important to understand the E-minimal model of W, Z. In general this is very difficult. In order to do this we obtain normal forms for the equations of X and Γ and we consider two cases. A general and a special one. In the general case we deform $\Gamma \subset X$ to a $\Gamma_0 \subset X_0$ having a simpler equation. Then we use the one parameter theory developed in the first part to show that in certain cases of interest, the divisorial contractions $E \subset Y \xrightarrow{f} \Gamma \subset X$ and $E_0 \subset Y_0 \xrightarrow{f_0} \Gamma_0 \subset X_0$ form a family. Then we explicitely construct the contraction f_0 and using results on deformations of terminal singularities we obtain information about f. In the special case we work directly with the equation of X. A complete classification of such contractions is given in Theorem 3.6.

This work has been done during my stay at the Max Planck Institute für Mathematik.

Finally I would like to thank the referree for his many usefull comments and for helping me make this paper readable.

1. *Notation and terminology*

Let X be a normal algebraic variety defined over the field of complex numbers $\mathbb{C}$, and D a Weil divisor on it.

D is called $\mathbb{Q}$-Cartier if there is an integer $m > 0$ such that mD is Cartier. We say that X is $\mathbb{Q}$-factorial iff every Weil divisor on X is $\mathbb{Q}$-Cartier.

Suppose that $K_X + \Delta$ is $\mathbb{Q}$-Cartier. Let $f\colon Y \longrightarrow X$ is a birational morphism from a normal variety Y and let $E_i \subset Y$ be the exceptional divisors. Then it is possible to write

$$K_Y + f_*^{-1}\Delta \equiv f^*(K_X + \Delta) + \sum_i a(E_i, X, \Delta)E_i.$$

We call $a(E, X, \Delta)$ the discrepancy of E with respect to (X, Δ). If $\Delta = 0$, then we denote $a(E, X, 0)$ by $a(E, X)$. The pair (X, Δ) is called canonical (resp. terminal) iff $a(E, X, \Delta) \geq 0$ (resp. $a(E, X, \Delta) > 0$ for all exceptional divisors E [**18**].

An f-exceptional divisor $E \subset Y$ is called f-crepant if $a(E, X, \Delta) = 0$. The birational morphism f is called crepant iff all f-exceptional divisors are f-crepant. Let $Z \subset X$ be a closed subscheme. Then we define the number of crepant divisors with center in Z by

$$e(X; Z) = \#\{E; a(E, X) = 0 \text{ and } \operatorname{center}_X E \subset Z\}.$$

By a slight abuse of notation we set $e(X; D) = e(X; \operatorname{supp}(D))$, for a Weil divisor D. If $Z = 0$ then we denote $e(X; 0)$ by $e(X)$. If X is canonical then it is known that $e(X; Z)$ is finite [**13**].

Definition 1.1. The D-minimal model of X is a projective morphism $f\colon Y \longrightarrow X$ such that Y is normal, the exceptional set of g has codimension at least 2, $D' = f_*^{-1}D$ is $\mathbb{Q}$-Cartier and f-ample over X. If it exists then it is unique.

It is known that the existence of the D-minimal model is equivalent to the finite generation of the sheaf of algebras $R(X, D) = \oplus_{m\geq 0}\mathcal{O}_X(mD)$ [**7**].

Many important constructions in birational geometry are related to D-minimal models. The flop is one of the most important ones. A flopping contraction is a projective birational morphism $f\colon Y \longrightarrow X$ to a normal variety X such that the exceptional set of f has codimension at least 2 in Y and K_Y is numerically f-trivial. If D_Y is a $\mathbb{Q}$-Cartier divisor on Y such that $-(K_Y + D_Y)$ is f-ample, then a projective morphism $f'\colon Y' \longrightarrow X$ such that the exceptional set of f' has codimension at least 2 and that $K_{Y'} + D_{Y'}$ is f'-ample is called the D-flop of f. In other words, it is the D-minimal model of X, where $D = f_*(D_Y)$. Its existence is fundamental in the context of the minimal model program.

If X is a 3-fold with canonical singularities, then $R(X, D)$ is finitely generated and the D-minimal model exists. The higher dimensional case is one of the most important open problems of higher dimensional birational geometry.

A $\mathbb{Q}$-factorialization of X is a projective birational morphism $f\colon Y \longrightarrow X$ such that Y is normal and $\mathbb{Q}$-factorial and f is an isomorphism in codimension 1. If X is a canonical 3-fold then a $\mathbb{Q}$-factorialization as above exists [**7**].

Let X be a canonical n-fold. A terminalization of X is a crepant projective birational morphism $f\colon Y \longrightarrow X$ such that Y has terminal only singularities. If $n = 3$ then a terminalization exists [**17**].

Let $f\colon Y \longrightarrow X$ be a birational morphism between normal varieties and let D be a Weil divisor in X. Then we denote the birational transform $f_*^{-1}D$ of D in Y by D^Y.

Let $\mathcal{X} \longrightarrow T$ be a flat family of algebraic varieties. A family of Weil divisors over T is a Weil divisor $D = \sum_i a_i D_i$ on $\mathcal{X}$ such that D_i is flat over T for all T.

2. *Families of D-minimal models*

In order to decide whether the minimal model program runs in families of canonical 3-folds, it is important to investigate if D-minimal models can be constructed in families. More precisely we want to study the following problem.

Problem: Let $X \longrightarrow T$ be a family of canonical 3-folds and let D be a family of divisors in X. It is known that the D_t-minimal model of X_t exists [**7**] for all $t \in T$. Do they form an algebraic family? i.e., Does there exists a morpism $f\colon Y \longrightarrow X$ such that $f_t\colon Y_t \longrightarrow X_t$ is the D_t minimal model of X_t?

In particular, we want to know if canonical flops exist in families. More precisely.

Problem: Let $Y/T \xrightarrow{f} X/T$ be a projective morphism between families of canonical 3-folds such that $Y_0 \xrightarrow{f_0} X_0$ is a flopping contraction. Does there exist a projective morphism $Y' \xrightarrow{f'} X$ such that $Y'_0 \xrightarrow{f'_0} X_0$ is the flop of f_0?

Let A be an f-ample divisor and $D = f_*(-A)$. Then Y' is the D-minimal model of X and therefore this problem is a special case of the problem concering families of D-minimal models.

From now on and unless otherwise said, T will denote a smooth curve. Then the total spaces X and Y are 4-folds. Recently Shokurov showed that canonical 4-fold log flips exist [**19**] and therefore the minimal model program works in dimension four. Let $f\colon Y/T \longrightarrow X/T$ be as above. Then the 4-fold flop of f exists. Suppose it is $f'\colon Y' \longrightarrow X$. However this does not necessarily mean that $Y'_0 \longrightarrow X_0$ is the flop of f_0. The reason is that even though f is a small contraction, it is possible that f_t is small for all $t \neq 0$, but f_0 is divisorial. The next example shows that this is indeed possible.

Example 2.1. Let $X_0 \subset \mathbb{C}^4$ be the 3-fold defined by the equation $xy+yu+xu = 0$, where $x,\ y,\ z,\ u$ are the coordinates in $\mathbb{C}^4$. It's singularities are cDV points and its singular locus is the line $L_0 : x = y = u = 0$. Therefore X_0 is canonical. Let $X \longrightarrow \mathbb{C}^1$ be the one parameter smoothing of X_0 given by $xy + yu + xu + ut = 0$, where t is the parameter. Let Z in X be the family of divisors given by $x = u = 0$, and let $f\colon Y \longrightarrow X$ be the blow up of X along Z. Then $f_t\colon Y_t \longrightarrow X_t$ is the blow up of X_t along the divisor Z_t given by $x = u = 0$. If $t \neq 0$, then X_t is smooth and therefore f_t is an isomorphism. On the other hand, X_0 is singular along $L_0 \subset Z_0$ and hence f_0 is divisorial contracting an irreducible divisor onto L_0.

The case of a family of terminal threefolds is particularly nice and it is possible to give a complete answer.

Theorem 2.2. *Let $X \longrightarrow T$ be a one dimensional family of terminal 3-folds. Let D be a family of Weil divisors in X over T. Then there exists a morphism $f\colon Y \longrightarrow X$, such that $f_t\colon Y_t \longrightarrow X_t$ is the D_t-minimal model of X_t for all t.*

Proof. First I claim that X has terminal singularities. This follows from the following results.

Theorem 2.3 [**9**]. *Let $f\colon X \longrightarrow T$ be a flat morphism from a germ of an algebraic vatiety to a germ of a smooth curve. Assume that the central fiber $X_0 = f^{-1}(P)$ has only canonical singularities. Then so has the total space X as well as any fiber X_t of f. Moreover, the pair (X, X_0) is canonical.*

It is also known [**16**] that if X_0 is terminal, then so is X and X_t. An immediate consequence of the above Theorem is that if $g\colon W \longrightarrow X$ is a resolution of X, then there is no g-exceptional crepant divisor with center in X_0. Hence all crepant exceptional divisors of X dominate T.

Now let $g\colon W \longrightarrow X$ be a resolution of X and let E_i be the exceptional divisors. Run a $(W, 0)$ MMP over X. Since X is terminal, we arrive at a $Y' \xrightarrow{g'} X$ over T, such that g' is an isomorphism in codimension 1 and Y' is $\mathbb{Q}$-factorial. Therefore $D' = {g'_*}^{-1} D$ is $\mathbb{Q}$-Cartier, and hence so is D'_t. To make it g'_t-ample, run a $(Y'_t, -\epsilon D'_t)$ MMP over X_t. By [**14**, Prop. 11.4, Theorem 11.10], and since X is terminal, this can be done in families. Therefore we obtain $Y \xrightarrow{g} X$ with the required properties. □

The case of families of canonical 3-folds is much more complicated. In general it is not possible to construct D-minimal models in families as shown by example 2.16. The number of crepant divisors of the members of the family is important as the next Theorem shows.

Theorem 2.4. *Let $X \xrightarrow{\sigma} T$ be a proper family of canonical threefolds over a smooth curve T. Let D in X be a family of divisors in X over T and let $0 \in T$ be a closed point. Then*

$$e(X_0) \geq e(X_t)$$

for t in a small neighborhood of $0 \in T$. Moreover,

(i) *If $e(X_0) = e(X_t)$ for all t, then after a finite base change there is a morphism $Y \xrightarrow{g} X$, with Y $\mathbb{Q}$-factorial and terminal, such that $Y_t \xrightarrow{g_t} X_t$ is a terminalization of X_t, for all $t \in T$.*

(ii) *If $e(X_0; D_0) = e(X_t; D_t)$ for all t, then there is a morphism $Y \xrightarrow{g} X$, such that $Y_t \xrightarrow{g_t} X_t$ is the D_t-minimal model of X_t.*

We are now able to get some information about the existence of flops in families.

Corollary 2.5. *Let $Y/T \xrightarrow{f} X/T$ be a morphism between one parameter families of canonical 3-folds. Suppose that $f_0\colon Y_0 \longrightarrow X_0$ is a flopping contraction. If $e(X_0) = e(X_t)$ for all t, then there is a morphism $Y'/T \xrightarrow{f'} X/T$ such that $f'_0\colon Y'_0 \longrightarrow X_0$ is the flop of f_0.*

Proof. Let D_Y be a divisor in Y such that $-D_Y$ is f-ample. Let $D = f_*(D_Y)$. Then $-D_{Y,t}$ is f_t-ample for all t and therefore Y_t is the $(-D_t)$-minimal model of X_t, and the flop of f_t is the D_t-minimal model of X_t. Since $e(X_0) = e(X_t)$ for all

t, then from Theorem 2.4 follows that there exist a morphism $f' \colon Y' \longrightarrow X$ such that $f'_t \colon Y'_t \longrightarrow X_t$ is the flop of f_t. □

Remark 2.6. The condition that $e(X_t)$ is constant in the family is probably too strong and the D-minimal models may form a family even though this condition does not hold. For example, with notation as in the previous Corollary, suppose that f_0 contracts a chain of rational curves to a canonical double point on X_0 but X_t is only terminal for $t \neq 0$. Then the the flop of f_0 can be constructed in families as Propostion 2.11 shows even though $e(X_0) > e(X_t) = 0$, for $t \neq 0$.

Proof of Theorem 2.4. The claimed inequality between the number of crepant divisors of the central and general fiber is independent of base change and so we will perform it when necessary. First I claim that up to a finite base change we can assume that $e(X_t) = e(X)$, for general $t \in T$. Indeed, let $g \colon W \longrightarrow X$ be a log resolution of X such that all the g-crepant divisors are smooth. Let E be a g-crepant divisor. From the previous theorem it follows that the center of E is not contained in X_0 and since there are finitely many crepant divisors, then after removing finitely many points from T we can assume that all the g-crepant divisors dominate T. By generic smoothness, E_t and W_t are smooth, for general t, and hence $g_t \colon W_t \longrightarrow X_t$ is a resolution of X_t. Then by adjunction it follows that E_t is g_t-crepant. The problem here is that E_t may have more than one connected components and so in general we only get that $e(X_t) \geq e(X)$. However, I claim that after a finite base change, E_t is irreducible and smooth, for all irreducible crepant exceptional divisors E of X, and therefore $e(X_t) = e(X)$. So, let E be an irreducible g-crepant divisor of X and let $E \xrightarrow{h} D \xrightarrow{\tau} T$ be the Stein factorization of $f = \tau \circ g \colon E \longrightarrow T$, where $D = \operatorname{Spec}(f_* \mathcal{O}_E)$. Then h has connected fibers and τ is finite. We now make a base change with $D \longrightarrow T$. So, let $X_D = X \times_T D$, $W_D = W \times_T D$ and $E_D = E \times_T D$. Note that by the previous theorem, X_D is also canonical since all the fibers $X_{D,d}$ are canonical. Also $W_{D,d}$ is smooth for general $d \in D$. Moreover, by the universal property of fiber product we see that there is an embedding $E \subset E_d$, and by construction, $E \longrightarrow D$ has connected fibers. Hence E_d is irreducible, smooth and crepant for $X_{D,d} = X_t$, for general $t = \tau(d)$. Since D may not be normal, make another base change with its normalization $\overline{D}$. Repeat this process for any crepant divisor F of $W_{\overline{D}}$ such that F_d is not irreducible. Since the crepant divisors are at most $e(X_t)$, this process ends with a family $X' \longrightarrow T'$, with T' smooth, such that there is a log resolution $f' \colon W' \longrightarrow X'$, such that if E is any f'-exceptional crepant divisor, E_t is smooth and irreducible and hence $e(X'_t) = e(X')$, for general t.

We may also assume that X is $\mathbb{Q}$-factorial. If this is not the case, then let $f \colon Y \longrightarrow X$ be a $\mathbb{Q}$-factorialization, which exists by the MMP in dimension 4. Then $e(X) = e(Y)$. Moreover, $Y_t \longrightarrow X_t$ is an isomorphism in codimension 1 for general t and hence $e(Y_t) = e(X_t) = e(X) = e(Y)$. The central fiber contraction $Y_0 \longrightarrow X_0$ may be divisorial, but in any case it is crepant and hence Y_0 is normal and canonical and $e(Y_0) \leq e(X_0)$. Hence in addition to our hypothesis we may also assume that X is $\mathbb{Q}$-factorial and $e(X_t) = e(X)$.

Let $n = e(X)$ and let $E_1, \dots, E_n$ be the crepant exceptional divisors of X. Then by standard MMP arguments, we may extract them from X with a series of crepant morphisms

$$Y = X_n \xrightarrow{f_n} X_{n-1} \xrightarrow{f_{n-1}} X_{n-2} \longrightarrow \cdots \longrightarrow X_1 \xrightarrow{f_1} X \tag{2.1}$$

where $f_i \colon X_i \longrightarrow X_{i-1}$ is crepant and its exceptional set is E_i. Then by Theorem 2.3 the centers of these divisors are not contained in X_0. Hence $X_{i,0}$ is irreducible and $E_i \cap X_{i,0}$ is a divisor. Moreover, I claim than $X_{i,0}$ is normal and canonical. Indeed, inductively it easily follows that $K_{X_i} + X_{i,0} = f_i^*(K_{X_{i-1}} + X_{i-1,0})$ and hence since (X, X_0) canonical, $(X_i, X_{i,0})$ is canonical as well and therefore $X_{i,0}$ is normal. Moreover, by adjunction it follows that $K_{X_{i,0}} = f_i^* K_{X_{i-1,0}}$ and hence $X_{i,0}$ is canonical and $E_i \cdot X_{i,0}$ is a crepant divisor for $X_{i-1,0}$ (which may be reducible). Therefore $e(X_0) \geq n = e(X_t)$.

Suppose now that $e(X_0) = e(X_t)$. Let $f \colon Y \longrightarrow X$ be the composition of the maps f_i in (2.1) above. Then by its construction, Y is a $\mathbb{Q}$-factorial terminal 4-fold and since $e(Y_t) = 0$ for general t, Y_t is terminal as well for general t. Moreover, by the above discussion it follows that Y_0 is irreducible and $e(Y_0) = 0$, and hence Y_0 is terminal too. Now $Y \longrightarrow X$ satisfies all the conditions of Theorem 2.4.1.

Now suppose that $D \subset X$ is a family of divisors such that $e(X_0; D_0) = e(X_t; D_t)$, for all t. Let $Z \xrightarrow{g} X$ be the D-minimal model of X, which exists by the MMP in dimension 4. Then $Z_t \longrightarrow X_t$ is an isomorphism in codimension 1 for general t and therefore $e(Z_t) = e(X_t)$. I now claim that $Z_0 \longrightarrow X_0$ is also an isomorphism in codimension 1 and hence it is also the D_0-minimal model of X_0. Suppose not. Let $D' = f_*^{-1} D$. Then by the definition of D-minimal models, $-D'$ is g-ample and hence if $Z_0 \longrightarrow X_0$ is divisorial, then the center of any g_0-exceptional divisor is contained in D_0. Therefore, $e(D'_0; Z_0) < e(D_0; X_0) = e(D_t; X_t) = e(D'_t; Z_t)$, which is impossible from the first part of the proof.

□

Remark. The condition $e(X_0) = e(X_t)$ is not sufficient for the existence of a morphism $g \colon Z \longrightarrow X$ such that $g_t \colon Z_t \longrightarrow X_t$ is a $\mathbb{Q}$-factorialization of X_t, for all t, as it was mistakenly claimed in [**22**]. The reason is that there may be divisors in X_0 that do not deform with X_0, i.e., they do not extend to a divisor in X. For example, let $X = (xy - zu + t = 0) \subset \mathbb{C}^4$. Then $X_0 = (t = 0)$ is the ordinary double point $xy - zu = 0$, and X_t for $t \neq 0$ is smooth. X_0 is not $\mathbb{Q}$-factorial and there is no morphism $g \colon Y \longrightarrow X$ such that $g_0 \colon Y_0 \longrightarrow X_0$ is a $\mathbb{Q}$-factorialization of X_0 because if there was such a morphism g, then g would be an isomorphism in codimension 1 which is impossible since X itself is smooth.

Unfortunately, the proof of Theorem 2.4 is using the existence of the MMP in dimension 4. It would be desirable to have a proof of it that does not involve the MMP, like in dimension 3, and constructive if possible. In general I do not know how to do this but in the next special case that is of interest in the study of divisorial contractions, it can be done. Moreover, in this case there is no restriction on the dimension of the base T.

Theorem 2.7. *Let $X \longrightarrow T$ be a family of index 1 canonical 3-folds, with T smooth. Let $Z \subset X$ be a smooth effective divisor in X flat over T. Moreover, assume that*

(i) *$X_t^{sing} \cap Z_t$ is a smooth curve L_t. Let $S_t = Spec\mathcal{O}_{X_t, L_t}$. This is a DuVal* *surface singularity defined over* $\mathbb{C}(t)$. $Z_t|_{S_t}$ *is a line through the origin such that the extended dual graph of* S_t *and* $Z_t|_{S_t}$ *is independent of* t. *This means that* S_t *are isomorphic singularities and that the position of* $Z_t|_{S_t}$ *in the fundamental cycle does not change with* t.

(ii) X_0 *is either cDV along* Z_0, *or there is a smooth curve* $L \subset X$ *that dominates* T *such that* $L \cap X_t$ *is a reduced point* P_t *and* $\mathrm{mult}_L X = 2, 3$ *or* ≥ 3. *This means that* $P_t \in X_t$ *is elliptic with constant* $k = \mathrm{mult}_L X$. *Moreover, the singularities of the weighted blow up* $B^w_{P_0} X_0$ *over* P_0 *on the birational transform* $\tilde{Z}_t$ *are finitely many cDV points, and* $w = (1,1,1,1)$, $(3,2,1,1,)$ *or* $(2,1,1,1)$ *if* $k = 1$, 2, *or* ≥ 3 *respectively.*

Then there is a birational morphism $Y \longrightarrow X$ *flat over* T, *such that* $Y_t \longrightarrow X_t$ *is the* Z_t*-minimal model of* X_t, *for all* t.

Remark 2.8. The complicated assumptions of the previous theorem easily imply that $e(X_0; Z_0) = e(X_t; Z_t)$, for all t, and therefore if $\dim T = 1$, Theorem 2.7 follows from Theorem 2.4. However, its proof does not use the MMP in dimension 4 and it is constructive.

The proof of Theorem 2.7 is given at the end of section 2. The idea is the following. By performing a series of well chosen blow ups we arrive at a crepant morphism $h\colon W/T \longrightarrow X/T$ suct that W_t is irreducible and that $(h_t)^{-1}_* Z_t$ is $\mathbb{Q}$-Cartier. We then run a relative MMP in families on W and arrive at a morphism $Y \longrightarrow X$ with the properties claimed in the Theorem.

The difficulty here is that in general W_t is only canonical and therefore we want to run a MMP in families of canonical 3-folds. For families of terminal 3-folds it is known to work [**14**]. The only operations of the MMP that is not known if they can be performed in families of canonical 3-folds is the flop and flip. Since we are always concerned with crepant morphisms, only the flop is needed in our case.

In what follows we obtain some information about cases when flops can be performed in families. We also study the behaviour of the property of being $\mathbb{Q}$-factorial in families of canonical 3-folds, and we also obtain a few more technical results that are needed for the proof of Theorem 2.7.

In the terminal case, the property of being $\mathbb{Q}$-factorial is stable under deformations as the next Theorem shows.

Theorem 2.9 [**14**, Theorem 12.1.10]. *Let* $X \xrightarrow{f} T$ *be a 1-dimensional family of terminal 3-folds, and let* $Z \subset X$ *be proper and flat over* T. *Let* D *be a divisor on* X *such that* $D_0 = D|_{X_0}$ *is* $\mathbb{Q}$*-Cartier along* Z_0. *Then there is a neighborhood* V *of* $0 \in T$ *such that* D *is* $\mathbb{Q}$*-Cartier along* $f^{-1}(V) \cap Z$.

In particular it follows that D_t is $\mathbb{Q}$-Cartier along Z_t for t in a neighborhood of $0 \in T$.

Unfortunately the previous result is no longer true for families of canonical 3-folds and one must be very careful when dealing with such families. The following example shows that it is possible that D_0 is $\mathbb{Q}$-Cartier, but D_t is not for all $t \neq 0$.

Example 2.10. We will now construct a one parameter family of canonical 3-folds $X \longrightarrow T$, a closed subscheme $Z \subset X$ proper and flat over T and a divisor

$D \subset X$, such that D_0 is $\mathbb{Q}$-Cartier along Z_0, but D_t is not $\mathbb{Q}$-Cartier along Z_t for all $t \neq 0$.

Start with the family given by $Y : x^2 + y^2 z + xz^5 + tu^3 + u^{10} = 0$, with t the parameter. Let $\Gamma \subset Y$ be given by $x = y = u = 0$. Now let $W = B_\Gamma Y \xrightarrow{g} Y$ be the blow up of Y along Γ. There are two exceptional divisors E and F. E is a $\mathbb{P}^1$ bundle over Γ, and F a $\mathbb{P}^2$ bundle over the line $x = y = z = u = 0$. Now let $X = B_E W \xrightarrow{h} W$ be the blow up of W along E. Let E^X, F^X be the birational transforms of E and F in X. Then, F^X is proper over T, and I claim that E_0^X is $\mathbb{Q}$-Cartier along F_0^X, but E_t^X is not $\mathbb{Q}$-Cartier for all $t \neq 0$.

For $t = 0$, Y_0 is given by $x^2 + y^2 z + xz^5 + u^{10} = 0$. Now explicit calculations of the blow up in exactly the same way as those in the proof of Theorem 3.6, show that $E_0 + 5F_0$ is Cartier and E_0^X is $\mathbb{Q}$-Cartier along F_0^X. Moreover, E_0 is not Cartier. Suppose it was. Then take a general fiber of g_0, $l \subset E_0$. This is rationally equivalent to a line $l' \subset F_0 = \mathbb{P}^2$. Then $-1 = E_0 \cdot l = E_0 \cdot l' > 0$.

Let us check now what happens in nearby fibers. For $t \neq 0$, Y_t is given by $x^2 + y^2 z + xz^5 + u^3 = 0$. Calculate the blow up of Γ now. In the affine chart $u \neq 0$ we have the coordinates $x = xu$, $y = yu$. Then W_0 is given by $x^2 u + y^2 zu + xz^5 + u^2 = 0$, and E_0, F_0 are given by $x = u = 0$ and $z = u = 0$ respectively. Now blow up E_0 to get X_0. In the affine chart $u = ux$, X_0 is given by $x^2 u + y^2 zu + z^5 + u^2 x = 0$, $E_0^{X_0}$ by $x = y^2 u + z^4 = 0$, $F_0^{X_0}$ by $z = u = 0$, and the h-exceptional divisor B by $x = z = 0$. Now it easy to see that $5F_0^{X_0}$ is Cartier and that X_0 is smooth away from $E_0^{X_0}$. But then, if $E_0^{X_0}$ was $\mathbb{Q}$-Cartier along $F_0^{X_0}$, then it would in fact be $\mathbb{Q}$-Cartier. But this would imply that E_0 is also $\mathbb{Q}$-Cartier as well which is not true.

We want to run a MMP in families of canonical 3-folds. To do this it is important to study the problem of whether canonical flops form families. This is true for terminal flops [**14**]. For canonical I do not know the answer in general but Theorem 2.4 suggests that in general it should be no. It is then very interesting to find conditions under which canonical flops can be constructed in families. Corollary 2.5 gives some conditions in this direction but they are most likely too strong. If the singularities that we are studying are double points then flops can be constructed in families as the next propistion shows.

Proposition 2.11. *Let $f \colon Y/T \longrightarrow X/T$ be a morphism between families of $n-$folds over T. Suppose that $f_0 \colon Y_0 \longrightarrow X_0$ is a flopping contraction and that X_0 has only hypersurface double points. Then the flop $f_0' \colon Y_0' \longrightarrow X_0$ of f_0 exists and there is a projective morphism $f' : Y' \longrightarrow X$, such that $f_0' : Y_0' \longrightarrow X_0$ is the flop of f_0.*

Proof. By using the Weierstrass preparation theorem we may write $0 \in X_0$ as

$$x^2 + f(y_1, \ldots, y_n) = 0.$$

Therefore, X can be written as

$$x^2 + f(y_1, \ldots, y_n, t_1, \ldots, t_m) = 0.$$

where $t_1, \ldots, t_m$ are local analytic coordinates of T. Now we proceed as in the 3-fold terminal case [**14**]. If $-D$ is f-ample, then to prove the existence of the flop for f it suffices to show that $R(X, D) = \oplus_d \mathcal{O}_X(df_* D)$ is finitely generated. X has an involution i. Then $D + i^* D = 0$. Since $R(X, -D)$ is finitely generated, then so is $R(X,D)$. Therefore there is a morphism $f'\colon Y' \longrightarrow X$ which is the flop of f. Moreover, $Y'=Y$, $f'=i\circ f$ and the involution is also in every fiber. Therefore the flop exists in families. □

Proposition 2.12. *Let $f\colon Y/T \longrightarrow X/T$ be a morphism between two families of canonical 3-folds over T. Suppose that $f_0\colon Y_0 \longrightarrow X_0$ is a flopping contraction. Suppose that there is a morphism $g\colon Z \longrightarrow Y$ over T, such that $Z_0 \longrightarrow Y_0$ is a $\mathbb{Q}$-factorial terminalization of X_0. Then there is a morphism $f'\colon Y' \longrightarrow X$, such that $f_0'\colon Y_0' \longrightarrow X_0$, is the flop of f_0.*

Note that the conditions of this proposition are not always satisfied as shown by example 2.16.

Proof. Let $f_0'\colon Y_0' \longrightarrow X_0$ be the flop of f_0. Let $g_0'\colon Z_0' \longrightarrow Y_0'$ be a $\mathbb{Q}$-factorial terminalization of Y_0'. Z_0, Z_0' are birational and nef over X_0. Therefore, $\phi_0\colon Z_0 \dashrightarrow Z_0'$ is an isomorphism in codimension 1 and a composition of terminal flops [**7**]. Terminal flops exist in families [**14**] and therefore ϕ_0 extends to a map $\phi\colon Z \dashrightarrow Z'$ over T. Then by [**15**], there is a morphism $g'\colon Z' \longrightarrow Y'$ over X, extending g_0'. Hence $f'\colon Y' \longrightarrow X$ is the required flop. □

The following lemmas are needed for the proof of Theorem 2.7.

Lemma 2.13. *Let $Z \subset X$ be a smooth divisor in a normal variety X. Suppose that X has canonical hypersurface singularities only, and that every irreducible component of $X^{sing}\cap Z$ is smooth. Let $Y = B_Z X \xrightarrow{f} X$ be the blow up of X along Z. Let $Z' = f_*^{-1}Z$, be the birational transform of Z in Y. Then*

(i) *Y is normal and has hypersurface singularities only.*
(ii) *$K_Y = f^*K_X$.*
(iii) *Every irreducible component E_i of the f-exceptional divisor E is smooth.*
(iv) *Every component of $Y^{sing}\cap Z'$ is smooth.*

Proof. Let $n=\dim X$. The results are all local around the singularities of X and therefore we may assume that $X \subset \mathbb{C}^{n+1}$. Let $f\colon B_Z\mathbb{C}^{n+1} \longrightarrow \mathbb{C}^{n+1}$ be the blow up of $\mathbb{C}^{n+1}$ along Z. Then $B_Z\mathbb{C}^{n+1}$ is smooth and $Y = B_Z X \subset B_Z\mathbb{C}^{n+1}$ has codimension 1 and hence Y has hypersurface singularities only. Let F be the f-exceptional divisor. Then by adjunction, $K_{B_Z\mathbb{C}^{n+1}} = f^*K_{\mathbb{C}^{n+1}} + F$, and $f^*X = Y + F$. Therefore

$$K_Y = (K_{B_Z\mathbb{C}^{n+1}} + Y)|_Y = f^*K_X.$$

Since $\mathbb{C}^{n+1}$ is smooth and X canonical, the proof of [**13**, Theorem 5.34] show that the pair $(\mathbb{C}^{n+1}, X)$ is also canonical. By adjunction we see that

$$K_{B_Z\mathbb{C}^{n+1}} + Y = f^*(K_{\mathbb{C}^{n+1}} + X).$$

Therefore the pair $(\mathbb{C}^{n+1}, Y)$ is also canonical. Hence by [**13**, Proposition 5.51], Y is normal. Finally, since every irreducible component of $X^{sing}\cap Z$ is smooth and $f^{-1}(z) = \mathbb{P}^1$, for all $z \in Z$, it follows that every irreducible component of $F \cap Y$ is also smooth.

Now let L be a component of the singular locus of X that lie on Z. Then this is either a point or a smooth curve. If it is a point then there is nothing to show.

So assume it is a smooth curve. At the generic point of L, X is a surface DuVal singularity, say X_L. Let $U \longrightarrow X_L$ be the minimal resolution and E_i, $i = 1, \ldots, n$ the exceptional curves. Z corresponds to a line through the singularity. We will now consider cases with respect to the type of the singularity X_L and the position of Z in the dual graph.

Case 1: X_L is an A_n singularity and Z intersects E_k in the dual graph.

Subcase 1: $(k, n) = (m+1, 2m+1)$, for some m. Then by [**20**], the f-exceptional divisor F is E_{m+1}, and hence again by [**20**], Z' is smooth over the generic point of L, and then $Z' \cap Y^{sing}$ over L is a finite set of points and perhaps a fiber of f which is a $\mathbb{P}^1$. Note that in this case it is possible that a component of Y^{sing} is a singular curve, but it does not lie on Z'.

Subcase 2: $(k, n) \neq (m+1, 2m+1)$, for any m. In this case, Y_L has two singular point, $P \in Z'$ and Q. Therefore $Y^{sing} \cap Z'$ consists of two curves over L and possibly some fibers of f. It is now clear that all these curves must be smooth.

Case 2: X_L is either D_n or E_i, $i = 6,\ 7,\ 8$ type DuVal singularity. These cases are treated exactly similar as the A_n cases and we will not do them. □

LEMMA 2.14. *Let* $(0 \in X)$ *be a canonical 3-fold singularity such that the general hyperplane section* H *of* X *through* 0 *is an elliptic surface singularity with invariant* $k = 1,\ 2$ *or* 3. *Let* $0 \in Z \subset X$ *be a smooth divisor. Let* $Y \xrightarrow{f} X$ *be either the blow up of* X *at* 0 *if* $k = 3$, *or the* $(3,2,1,1)$ *or* $(2,1,1,1)$ *weighted blow up of* X *at* 0 *in the case that* $k = 2$ *or* 1. *Then the birational transform* Z' *of* Z *in* Y *is smooth and* $Y^{sing} \cap Z$ *is a smooth rational curve or a point or empty.*

Proof. Suppose that $k = 3$. Then Y is just the blow up of X at 0 and therefore $Z' = B_0 Z$ is smooth.

Now suppose that $k = 2$ or 1. Then $\mathrm{mult}_0 X = 2$ [**13**, Theorem 4.57]. Let Z be given by $x = y = 0$. Then X is given by an equation of the form

$$xf(x,y,z,t) + yg(x,y,z,t) = 0.$$

Since $t = 0$ is elliptic, we can write the equation of X as

$$x^2 + xf(x,y,z,t) + yg(x,y,z,t) = 0.$$

By using the Weierstrass preparation theorem, the equation of X can be written as

$$x^2 + 2x\phi(y,z,t) + yh(y,z,t) = 0. \tag{2.1}$$

Moreover, $\phi(y,z,t),\ h(y,z,t) \in (y,z,t)^2$, because otherwise $0 \in X$ is cA_n. Eliminate x from (1). Then the equation of X becomes

$$F(x,y,z,t) = x^2 + yh(y,z,t) - \phi^2(y,z,t) = 0, \tag{2.2}$$

and Z is given by $x - \phi(y,z,t) = y = 0$. Moreover from the previous discussion, $\phi(y,z,t)$, $h(y,z,t) \in (y,z,t)^2$. Now consider cases with respect to h.

Case 1: $\mathrm{mult}_0 h(y,z,t) \geq 3$. Then assign weights to x, y, z and t as follows. Let $w(x) = 2$ and $w(y) = w(z) = w(t) = 1$. Let $Y = B_0^w X \xrightarrow{f} X$ be the $(2,1,1,1)$ weighted blow up of X. We want to understand the birational transform Z' of Z in Y.

Claim: $Z' = B_0 Z$ and therefore Z is smooth. By the definition of the weighted blow up,

$$Z' = \operatorname{Proj} \oplus_{n\geq 0} \frac{m^w(n)+I_Z}{I_Z},$$

and $I_Z = (x-\phi(y,z,t), y)$. We may look at Z as given by $x - \phi_1(z,t) = 0$ in $\mathbb{C}^3_{x,z,t}$, where $\phi_1(z,t) = \phi(0,z,t)$. Note that $\phi_1(z,t) \neq 0$, since otherwise $\phi(y,z,t) = y\lambda(y,z,t)$ and hence $2Z$ given by $x^2 = y = 0$ will be Cartier. Then $\text{mult}_0\phi_1(z,t) \geq 2$. Now

$$\frac{m^w(1)+I_Z}{I_Z} = (\overline{x},\overline{z},\overline{t}) = (\overline{z},\overline{t},\phi_1(\overline{z},\overline{t})) = (\overline{z},\overline{t}) = \frac{m+I_Z}{I_Z}$$

In general,

$$\frac{m^w(n)+I_Z}{I_Z} = (\overline{x}^i\overline{t}^j\overline{z}^k; 2i+j+k\geq n) =$$
$$(\phi_1(\overline{z},\overline{t})^i\overline{t}^j\overline{z}^k; 2i+j+k\geq n) = (\overline{t}^j\overline{z}^k; j+k\geq n) = \frac{m^n+I_Z}{I_Z}$$

Therefore, $Z' = B_0Z$, and hence Z' is smooth. Note that this is the case that $k=2$.

Case 2: $\text{mult}_0 h(y,z,t) = 2$, and $k=1$. Let $h_2(y,z,t)$ be the degree 2 part of $h(y,z,t)$. Since the section $t=0$ is elliptic, the cubic term of $F(x,y,z,0)$ must be a cube because otherwise the section $t=0$ is DuVal. Therefore,

$$h_2(y,z,t) = ay^2 + tl(y,z,t),$$

where $l(y,z,t)$ is linear in y, z, t. Therefore the equation of X becomes

$$x^2 + ay^3 + ytl(y,z,t) + yh_{\geq 4}(y,z,t) - \phi^2(y,z,t) = 0. \tag{2.3}$$

If $l(y,z,t)\neq 0$, then one of y^2z, yzt, yt^2, appears in the above equation and the section $z=t$ is given by $x^2+\psi(y,t)=0$, and $\psi_3(y,t)$ is not a cube and hence it must be DuVal. Therefore $l(y,z,t)=0$. Therefore the equation of X becomes

$$x^2 + y^3 + yh_{\geq 4}(y,z,t) - \phi^2(y,z,t) = 0, \tag{2.4}$$

where $\text{mult}_0\phi \geq 2$. Moreover, Z is given by $x-\phi(y,z,t) = y = 0$, and as before, $\phi(0,z,t)\neq 0$. By using the Weierstrass preparation theorem it is possible to write

$$x^2 + y^3 + yh_{\geq 4}(y,z,t) - \phi^2(y,z,t) =$$
$$x^2 + u_1[y^3 + \alpha(z,t)y^2 + \beta(z,t)y + \gamma(z,t)] =$$
$$u_1[(\frac{x}{\sqrt{u_1}})^2 + y^3 + \alpha(z,t)y^2 + \beta(z,t)y + \gamma(z,t)]$$

From the above equations, it is clear that $\gamma(z,t) = unit\cdot\phi^2_{\geq 2}(0,z,t) = -\psi^2_{\geq 2}(z,t)$. Therefore Z is given by $x - \psi_{\geq 2}(z,t) = y = 0$. Now eliminate y^2. The equation of X can be written

$$x^2 + y^3 + 3\alpha(z,t)y^2 + \beta(z,t)y - \psi^2_{\geq 2}(z,t) = 0.$$

The change of variables $y\mapsto y-\alpha(z,t)$ makes the equation of X

$$x^2 + y^3 + (y-\alpha(z,t))\delta(z,t) - \alpha^3(z,t) - \psi^2(z,t) = 0, \tag{2.5}$$

where $\delta(z,t) = \beta(z,t) - 3\alpha^2(z,t)$. Z is given by $x-\psi_{\geq 2}(z,t) = y-\alpha(z,t) = 0$. Now it is easy to see that $\text{mult}_0(\alpha^3(z,t)+\psi^2(z,t)) \geq 6$. Therefore, $\text{mult}_0(\psi)\geq 3$ and $\text{mult}_0(\alpha)\geq 2$. Now an argument as in case 1 shows that $Z' = B_0Z$ and therefore it is smooth.

Finally it remains to justify the statement about the singular locus of Y. The singularities of Y on Z' over 0 are on the intersection of Z' and the exceptional divisors. But since f is a certain weighted blow up, this intersection is a smooth rational curve. □

The next elementary result is needed for the proof of Theorem 2.7.

Lemma 2.15. *Let T be a smooth curve and $Y/T \xrightarrow{f} X/T$ be a morphism over T. Assume that X_0 is a canonical 3-fold. Suppose that $K_Y = f^*K_X$. Then $\dim f^{-1}(x) \leq 2$ for any $x \in X_0$.*

Proof. As before we see that the pair (X, X_0) is canonical. Therefore, for any $x_0 \in X_0$, $x_0 \in X$ is a terminal singularity. If there is a divisor $E \subset f^{-1}(x_0)$, then E is crepant over x_0 which is impossible. Therefore $\dim f^{-1}(x_0) \leq 2$. □

2.1. *Proof of Theorem 2.7*

Suppose that X_0 is given by $F(x,y,z,w) = 0$ in $\mathbb{C}^4$. Then X is given by $F(x,y,z,w) + t\Psi(x,y,z,w) = 0$ in $\mathbb{C}^5$. Let $X_1 \xrightarrow{f_1} X$ be the $(1,1,1,1,1)$, or $(2,1,1,1,1)$, or $(3,2,1,1,1)$ weighted blow up of X along L. Consider the $(1,1,1,1,1)$ blow up case first. Let $\tilde{\mathbb{C}} = B_0\mathbb{C}^5$. Then

$$K_{\tilde{\mathbb{C}}} = f_1^*K_{\mathbb{C}^5} + 3E$$
$$f_1^*X = Y + 3E.$$

Therefore

$$K_{X_1} = f_1^*K_X.$$

By lemma 2.15, it follows that $\dim f_1^{-1}(x) \leq 2$ for any $x \in X_0$. Therefore $X_1 \longrightarrow T$ is also a family of canonical 3-folds. Moreover, $e(X_1) < e(X)$. By lemma 2.14, $Z_1 = (f_1)_*^{-1}Z$ is smooth and $X_1^{sing} \cap (Y_1)_0$ is a union of smooth rational curves. Let $X_2 \xrightarrow{f_2} X_1$ be the blow up of X_1 along Z_1. Then by lemma 2.13, $K_{X_2} = f_2^*K_{X_1}$, X_2 is normal and $e(X_2) < e(X_1)$. Moreover, the conditions of part 1 of the theorem guarantee that there is a smooth f_2-exceptional divisor (over the generic point of L_0 it is just the blow up of a line through a DuVal singularity). Moreover, again by lemma 2.13, every component of the singular locus of X_2 that lies on Z_2 is smooth. If it is just an isolated set of points then stop. If not, then blow up the f_2-exceptional divisor that they lie on, which as mentioned is smooth, to get a crepant morphism $X_3 \xrightarrow{f_3} X_2$, with $e(X_3) < e(X_2)$. Continue this process of blowing up exceptional divisors to get a sequence

$$\cdots \longrightarrow X_k \xrightarrow{f_k} X_{k-1} \longrightarrow \cdots \xrightarrow{f_3} X_2 \xrightarrow{f_2} X_1 \xrightarrow{f_1} X,$$

such that

$$e(X_1) > e(X_2) > \cdots e(X_k) > \cdots$$

Since there are finitely many crepant divisors, this process must stop and there is an $n > 0$ such that either $e(X_n) = 0$ and hence X_n is terminal, or by lemma 2.13 there are finitely many only singularities on Z_n. Moreover, $g_n \colon X_n \longrightarrow X$ is crepant and there are finitely many singularities only on every g_n-exceptional curve. Let E_i be the g_n-exceptional divisors which by lemma 2.13 are all smooth. As we mentioned above, $(X_n)_0$ has only isolated hypersurface singularities on Z_n. Therefore we can obtain a $\mathbb{Q}$-factorialization of it, W, by just blowing up Z_n and the exceptional E_i. Now consider the map $W/T \longrightarrow X/T$. A $\mathbb{Q}$-factorialization of Z_0 can be obtained by running a $(W_0, \epsilon \sum E_i^W)$ MMP over X_0. Since W_0 has finitely many terminal singularities over any point of X, then by Propositions 2.11 and 2.12, the operations of the MMP extend in the family. Therefore we obtain a morphism $Y/T \xrightarrow{f} X/T$, such that $Y_0 \xrightarrow{f_0} X_0$ is a $\mathbb{Q}$-factorialization of Z_0. We now want to conclude that $Y_t \xrightarrow{f_t} X_t$ is a $\mathbb{Q}$-factorialization of Z_t for t in a neighborhood of 0. As we have seen in example 2.10, this is not automatic in a family of canonical threefolds. However, the conditions of theorem 2.7 mean that $e(X_0) = e(X_t)$, for all t. Since $Y_0 \longrightarrow X_0$ is small, then by semicontinouity of fiber dimensions it follows that $Y_t \longrightarrow X_t$ is also small in a neighborhood of 0. Therefore, $e(Y_0) = e(Y_t)$. But this means that when we run the $(W_0, \epsilon \sum E_i^W)$ MMP, it is not possible that there is a divisorial contraction in the central fiber and small away from it which must be the case if Y_t fails to be $\mathbb{Q}$-factorial. The theorem now follows. □

The next example shows that the conditions of Theorem 2.7 are needed in order for the D-minimal models to form a family.

Example 2.16. In this example we construct a one parameter family of canonical 3-folds $Y \longrightarrow \mathbb{C}^1$ and a family of Weil divisors E in Y, such that the E_u-minimal models of Y_u do not form a family, where u is the parameter.

Let $X_0 \subset \mathbb{C}^4$ be the 3-fold terminal singularity given by $xy + z^3 + t^3 = 0$, where $x,\ y,\ z,\ t$ are the coordinates of $\mathbb{C}^4$. Let $X \longrightarrow \mathbb{C}^1$ be a one parameter deformation of X_0 given by $uz^2 + xy + z^3 + t^3 = 0$, where u is the parameter. X_0 has a cA_2 singularity but X_t only cA_1, for $t \neq 0$. Let $\Gamma \subset X$ be the plane given by $x = z = t = 0$ and let $f\colon Y = B_\Gamma X \longrightarrow X$ be the blow up of X along Γ. Then an explicit calculation of Y shows that the exceptional set of f consists of two divisors. A $\mathbb{P}^1$-bundle E over Γ, and a $\mathbb{P}^2$-bundle F over the line $x = y = z = t = 0$. Moreover, E_u and F_u are not $\mathbb{Q}$-Cartier for all u.

Claim: The E_u-minimal models of Y_u do not form a family. Suppose they do. Then there is a morphism $g\colon Z \longrightarrow Y$ such that $g_u\colon Z_u \longrightarrow Y_u$ is the E_u-minimal model of Y_u. Hence E_u^Z and F_u^Z are $\mathbb{Q}$-Cartier. Now we can contract F_0^Z to a terminal singularity. Let $p_0\colon Z_0 \longrightarrow W_0$ be the contraction. Then the whole process is described by the following diagram:

$$\begin{tikzcd}
 & Z_0 \arrow[dl, "g_0"'] \arrow[dr, "p_0"] & \\
Y_0 \arrow[dr, "f_0"'] & & W_0 \arrow[dl, "q_0"] \\
 & X_0 &
\end{tikzcd}$$

where $q_0\colon W_0 \longrightarrow X_0$ contracts E_0^W onto the curve $\Gamma_0 :\ x = z = t = 0$. If the E_u-minimal models of Y_u form a family and a morphism $g\colon Z \longrightarrow Y$ exists as above, then p_0 and q_0 extend to the family [**14**, Proposition 11.4], and the above diagram is a specialization of the diagram

$$\begin{tikzcd} & Z \arrow[dl, "g"'] \arrow[dr, "p"] & \\ Y \arrow[dr, "f"'] & & W \arrow[dl, "q"] \\ & X & \end{tikzcd}$$

where W is a one parameter deformation of W_0. In particular there is a family of divisorial contractions $q\colon W \longrightarrow X$ such that for all u, $q_u\colon W_u \longrightarrow X_u$ is a divisorial contraction contracting an irreducible divisor E_u^W onto the line $\Gamma_u \subset X_u$ given by $x = z = t = 0$. However,

Claim: W_0 is terminal of index 3, but W_u is terminal of index 2, for all $u \neq 0$. Since the index is constant in families of terminal 3-folds, we get a contradiction and therefore the E_u-minimal models of Y_u do not form a family.

We now proceed to justify all the previous claims. By construction, $Y_u = B_{\Gamma_u}X_u$. In the affine chart given by $x = xt$, $z = zt$, Y is given by

$$uz^2t + xy + z^3t^2 + t^2 = 0.$$

Set $t = 0$ to see that $f^{-1}(\Gamma) = E + F$, where E is given by $x = t = 0$ and F by $y = t = 0$. The other charts are checked similarly.

Assuming that the E_u-minimal models of Y_u form a family, the only thing that needs to be proved is the claim about the indices of W_u. X_0 is given by $xy+z^3+t^3 = 0$ and Γ_0 by $x = z = t = 0$. The general hyperplane section of X_0 through Γ_0 has an A_2 DuVal singularity. For $u \neq 0$, X_u is given by $uz^2 + xy + z^3 + t^3 = 0$. In this case, the general hyperplane section through Γ_u has an A_1 DuVal singularity. Therefore, W_u has index 2 if $u \neq 0$ [**20**, Theorem 5.1].

We will construct the contraction $W_0 \longrightarrow X_0$ explicitely. Y_0 is given by $xy + z^3t^2 + t^2 = 0$. E_0 is given by $x = t = 0$ and F_0 by $y = t = 0$. We want to construct the E_0-minimal model of Y_0, Z_0, and then contract F_0 to obtain W_0. Y_0 is given by $xy + (z^3 + 1)t^2 = 0$ and a straightforward calculation shows that its singular locus is the line $x = y = t = 0$. Moreover, $2E_0$ is Cartier at all points except at the three points given by $x = y = t = z^3 + 1 = 0$. Therefore, in order to make E_u $\mathbb{Q}$-Cartier all we have to do is blow up the ideal $I = (x, t^2)$ which defines $2E_0$. Let $g\colon Z_0 \longrightarrow Y_0$ be the blow up of I. Then g is an isomorphism in codimension 1 and moreover I claim that the g-exceptional curves C, are not contained in the birational transform F_0^Z and therefore $F_0^Z = F_0 = \mathbb{P}^2$. Then F_0^Z can indeed be contracted to a point by a K_{Z_0}-negative extremal contraction as claimed earlier. To see this describe Z_0 explicitely. By its construction, $Z_0 \subset \mathbb{C}^4 \times \mathbb{P}^1_{u,v}$ and is given by

$$\begin{gathered} xu - t^2v = 0 \\ xy + z^3t^2 + t^2 = 0 \end{gathered}$$

In the affine chart $v = 1$, Z_0 is given by

$$\begin{gathered} xu - t^2 = 0 \\ y + z^3u + u = 0 \end{gathered}$$

The g-exceptional curves are given by $x = y = t = z^3 + 1 = 0$, and $F_0^{Z_0}$ by

$y = t = u = 0$. Therefore no g-exceptional curve is contained in $F_0^{Z_0}$ and hence $F_0^{Z_0} = F_0 = \mathbb{P}^2$. Let $l \subset F_0^Z$ be a line. Then $l \cdot K_{Z_0} = l \cdot K_{Y_0} = -1 < 0$ and therefore $Z_0 \xrightarrow{p_0} W_0$ is a K_{Z_0}-negative extremal contraction as claimed.

We now proceed to find the index of W_0. We know that $2E_0^{Z_0}$ and $2F_0^{Z_0}$ are Cartier. Moreover there is a rational number $a > 0$ such that

$$K_{Z_0} = p^* K_{W_0} + aF_0^{Z_0}. \tag{2.6}$$

Let $l \subset F_0^{Z_0} = \mathbb{P}^2$ be a line. Then $l \cdot F_0^{Z_0} = -b/2$, where b is a positive integer. We want to find b.

$$l \cdot (E_0^{Z_0} + F_0^{Z_0}) = l \cdot g_0^*(E_0 + F_0) = l \cdot (E_0 + F_0) = -1.$$

Now, $l \cdot E_0^{Z_0} = 1/2(2E_0^{Z_0}) \cdot l$. This can easily be computed. $2E_0^{Z_0}$ is given by $x = t^2 = y + z^3u + u = 0$ and $F_0^{Z_0}$ by $y = t = u = 0$. We can take any line in $F_0^{Z_0}$, so take the one given by $y = t = z = u = 0$. The intersection $l \cap 2E_0^{Z_0}$ is a reduced point and therefore $l \cdot 2E_0^{Z_0} = 1$. Hence

$$l \cdot F_0^{Z_0} = -1 - l \cdot E_0^{Z_0} = -3/2.$$

Intersecting (6) with l we see that $a = 2/3$. Therefore the index of W_0 is 3.

3. *Applications to divisorial contractions.*

In this section we obtain a complete classification of terminal 3-fold divisorial contractions $f\colon E \subset Y \longrightarrow \Gamma \subset X$, in the case that Γ is a smooth curve and the general hyperplane section S of X through Γ has a D_5 DuVal singularity.

Definition 3.1. A divisorial contraction is a proper morphism $f : E \subset Y \longrightarrow \Gamma \subset X$, such that Y is $\mathbb{Q}$-factorial, $Y - E \cong X - \Gamma$, E is an irreducible divisor and $-K_Y$ is f-ample.

The contraction is called terminal if both X and Y have terminal singularities. We also need the following definition.

Definition 3.2 [**20**]. Let $0 \in S$ be a D_n DuVal singularity, and let $0 \in \Gamma \subset S$ be a smooth curve through the singularity. Let $U \xrightarrow{f} S$ be the minimal resolution and $\Gamma' = f_*^{-1}\Gamma$. Let



.

be the dual graph. Then Γ' intersects either E_1, or E_{n-1} or E_n. In the first case we call Γ of type DF_l, and in the others of type DF_r.

Construction of contractions 3.3. Let $\Gamma \subset X$ be a curve in a 3-fold X having at most index 1 terminal singularities. Suppose that the general hyperplane section S of X through Γ is DuVal and that the curve has at worst plane singularities. Then there is a divisorial contraction $f\colon Y \longrightarrow X$ contracting an irreducible divisor E onto Γ such that Y has at worst canonical singularities. Moreover such contraction is unique and can be obtained from the following diagram [**20**].

$$\begin{tikzcd}
Z \arrow[rr, dashed, "\phi"] \arrow[d, "h"'] & & Z' \arrow[d, "h'"] \\
W \arrow[dr, "g"'] & & Y \arrow[dl, "f"] \\
& X &
\end{tikzcd}$$

W is the blow up of X along Γ. There are two g-exceptional divisors. A ruled surface E over Γ, and a $F = \mathbb{P}^2$ over the singularity. Z is the E-minimal model of W. After a sequence of flips, F can be contracted by a birational morphism $h' \colon Z' \longrightarrow Y$, and we obtain the required contraction.

In order to understand when a terminal contraction exists, i.e., when Y is terminal, it is nececssary to study the singularities of Z and Z'. In general, calculating D-minimal models directly is difficult and the possible appearance of flips makes things worse. To overcome this difficulty we will use the one parameter theory developed in the first part of the paper to degenerate $\Gamma \subset X$ to a $\Gamma_0 \subset X_0$ having simpler equation and therefore easier to manipulate. Then we want to deform Y to Y_0. In order to do this we must study the following problem.

Problem: Let $X \longrightarrow T$ be a one parameter family of terminal 3-folds and let $\Gamma \subset X$ be flat over T, such that $\Gamma_t \subset X_t$ is a smooth curve for all t. Does there exist a morphism $f \colon Y \longrightarrow X$ such that $f_t \colon Y_t \longrightarrow X_t$ is a divisorial contraction contracting an irreducible divisor E_t onto Γ_t?

In general the answer is no. However under certain conditions they do form families, as the next result shows.

Corollary 3.4. *Let $X \longrightarrow T$ be a 1-dimensional family of terminal 3-fold singularities $P_t \in X_t$. Let $\Gamma \subset X$ be flat over T such that $\Gamma_t \subset X_t$ is a smooth curve through P_t. Let H_t be the general hyperplane section of X_t through Γ_t. Suppose it is an A_n DuVal singularity for all t and that Γ_t intersects an end of the fundamental cycle of H_t for all t. Then there exist a morphism $Y \xrightarrow{f} X$, such that $Y_t \xrightarrow{f_t} X_t$ is a terminal contraction contracting an irreducible divisor E_t onto Γ_t.*

Proof. Let $g \colon W \longrightarrow X$ be the blow up of X along Γ. Then as mentioned earlier, there are two g-exceptional divisors E and F. Moreover, from [**20**] it follows that W_t is *cDV* for all t. Moreover, the singular locus of W_t is the line $L_t = E_t \cap F_t$, and W_t is an A_{n-1} DuVal singularity at the generic point of L_t. Therefore the conditions of theorem 2.7 are satisfied and hence there is a morphism $h \colon Z \longrightarrow W$, such that $Z_0 \longrightarrow W_0$ is the E_0-minimal model of W_0. Now the construction described above can be done in families and hence there is a morphism $f \colon Y \longrightarrow X$ with the properties claimed. □

Note that the conditions of Corollary 3.4 are necessary as example 2.16 shows.

We can use the previous result to improve the result of [**20**, Theorem 5.1].

COROLLARY 3.5. *Let $E \subset Y \longrightarrow \Gamma \subset X$ be a terminal 3-fold divisorial contraction contracting an irreducible divisor E onto a smooth curve Γ. Suppose that the general hyperplane section S of X through Γ is A_n, and that Γ intersects an end of the fundamental cycle of S. Then Y has index $n+1$.*

Proof. By [**20**] it is possible to write the equation of X as

$$xy + z^{n+1} + t\phi(x,y,z,t) = 0,$$

where the curve Γ is given by $x = z = t = 0$. By a result of Hironaka-Rossi, the equation of X is equivalent to

$$xy + z^{n+1} + t\phi(x,y,z,t) + t^m = 0,$$

for $m >> 1$. Now we can deform X to X_0 given by $xy+z^{n+1}+t^m = 0$, and it can be explicitely seen with the same method as in example 2.16, that there is a terminal contraction $E_0 \subset Y_0 \longrightarrow \Gamma_0 \subset X_0$, and Y_0 has index $n+1$. These contractions form a family by Corollary 3.4. The index is constant in families of terminal 3-folds and hence the corollary follows. □

THEOREM 3.6. *Let $0 \in \Gamma \subset S \subset X$. Suppose that $P \in S$ is a D_5 singularity for the general S through Γ. Then*

(i) *If $\Gamma \subset S$ is of type DF_l, then there is no terminal contraction.*
(ii) *If $\Gamma \subset S$ is of type DF_r, then*
  (a) *If $0 \in X$ is a cD_4 singularity, then there is a terminal contraction.*
  (b) *If $0 \in X$ is a cD_5 singularity, then it is always possible to write the equation of $\Gamma \subset S \subset X$ as*

$$x^2 + y^2z + xz^2 + t[xz\psi(z,t) + axt^k + \phi_{\geq 3}(y,z,t)] = 0,$$

so that z^ν does not appear in $\phi_{\geq 3}(y,z,t)$ for any ν, $k \geq 1$ and $I_\Gamma = (x,y,t)$. Let $a_{i,j,k}$ denote the coefficient of $y^iz^jt^k$ in $\phi_{\geq 3}(y,z,t)$ and b the coefficient of xt^2 in the above equation (i.e., either 0 or a). Then a terminal contraction exists unless

(1)

$$a_{0,0,4} = a_{1,0,2} = 2a_{0,1,2} - b\psi(0,0) = 4a_{0,0,3} - b^2 = 0$$

or

(2)

$$\begin{aligned} a_{0,2,1}^2 - ba_{0,2,1} + a_{0,0,3} &= 0 \\ a_{0,1,2} - a_{0,2,1}\psi(0,0) &= 0. \end{aligned}$$

If there is a terminal contraction $E \subset Y \xrightarrow{f} \Gamma \subset X$, then Y has index 4.

EXAMPLE 3.7. Let $0 \in \Gamma \subset S \subset X$ be given by

$$X : x^2 + y^2z + 2xz^2 + t\phi_{\geq 4}(y,z,t) = 0,$$

and Γ by $x = y = t = 0$. Then there is no terminal divisorial contraction $Y \longrightarrow X$ contracting an irreducible suface onto Γ.

COROLLARY 3.8. *Let $\Gamma \subset X$ as in the theorem. Let $E \subset Y \xrightarrow{f} \Gamma \subset X$ the* *canonical divisorial contraction, which by [**20**] always exists. Let Σ be the general section of X through* 0*. Then Y is terminal iff $\Sigma' = f_*^{-1}\Sigma$ is normal.*

The corollary follows immediately from the proof of Theorem 3.6. At this point I would like to point out that the conclusion of the corollary is still true in the case that the general hyperplane section S of X through Γ is a D_{2n} DuVal singularity as well. However it is not true in general. In particular there are examples when $0 \in X$ is a cA_2 singularity and Σ' is not normal [**21**].

Proof Proof of Theorem 3.6. The method that we are going to use is based on the method that appears in [**20**] which was described in the beginning of this section. Fix notation as in 3.3.

The idea of the proof is the following. First we obtain normal forms for the equations of Γ and X. Then we distinguish two cases with respect to the equations of $\Gamma \subset X$. A general one and a special one, and we treat them seperately. In the general case, we degenerate $\Gamma \subset X$ to a $\Gamma_0 \subset X_0$ whose equation is simpler. Then we follow the method described in 3.3 to construct the contraction $Y_0 \longrightarrow X_0$. We then use the one parameter theory developed in the first part of the paper to show that the contractions form a family. Now a deformation of a terminal singularity is again terminal [**16**] and hence we obtain information about the singularities of Y from what we know about Y_0.

In the special case we work explicitely with the equation of X and Γ. The main difficulty is to describe Z and the possible existence of flips. We show that Z can be obtained from the following diagram:

$$\begin{tikzcd}
 & W_2 \arrow[dl, "h_2"'] \arrow[dr, "h'"] & \\
W_1 \arrow[dr, "h_1"'] & & Z \arrow[dl, "h"] \\
 & W &
\end{tikzcd}$$

where W_1 is the blow up of W along E. Let F_1 be the h_1-exceptional divisor. Then by blowing up a suitable multiple of E^{W_1} we make it $\mathbb{Q}$-Cartier and get the E^{W_1}-minimal model of W_1 which we call W_2. We then show that no h_2-exceptional curve is contained in $F_1^{W_2}$ and hence we can contract $F_1^{W_2}$ to obtain Z. Moreover from the above construction follows that no h-exceptional curve is contained in F^Z, and therefore $F^Z \cong F \cong \mathbb{P}^2$. Hence F^Z can be contracted to a terminal singularity and therefore no flips exist in the construction of 3.3. Then in order to decide whether Y is terminal or only canonical we need to study the singularities of Z away from F^Z. If they are isolated terminal, then so is Y. If not, then Y is only canonical.

Part 1. of the theorem follows from [**20**, Theorem 6.1]. We now proceed to justify all the steps described above. By [**20**, Proposition 4.8], in suitable analytic coordinates, the equation of X is

$$x^2 + y^2z + xz^2 + t[xz\psi(z,t) + axt^k + \phi_{\geq 2}(y,z,t)] = 0 \tag{3.1}$$

with $k \geq 1$.Moreover, Γ is given by $x = y = t = 0$, and the monomials y^2, yz and z^ν do not appear in $\phi_{\geq 2}(y,z,t)$, for any ν.

The appearance of the monomials yt and t^2 in $\phi_{\geq 2}(y,z,t)$ complicate the calculations a lot and it is best to consider two cases with respect to these.

Case 1. One or both of the monomials yt and t^2 appear in $\phi_2(y,z,t)$. Suppose that yt exists. Then write the equation of X as

$$x^2+y^2z+xz^2+yt^2+t\psi(x,y,z,t)=0.$$

We can now deform X to X_0 given by

$$x^2+y^2z+xz^2+yt^2=0.$$

Let $\mathbf{X}\longrightarrow\Delta$ be the deformation over the unit disk. Let $\mathbf{\Gamma}\subset\mathbf{X}$ be the deformation of Γ given by $x=y=t=0$, and let $\mathbf{W}\xrightarrow{\mathbf{g}}\mathbf{X}$ be the blow up of $\mathbf{X}$ along $\mathbf{\Gamma}$. Let $\mathbf{E}$, $\mathbf{F}$ be the two $\mathbf{g}$-exceptional divisors. Then $\mathbf{W}_u\cong W=B_\Gamma X$ for $u\neq 0$ in Δ, and $\mathbf{W}_0\cong W_0=B_{\Gamma_0}X_0$. Now one can check that the family $\mathbf{W}$ satisfies the conditions of theorem 2.7. Therefore there exists a morphism $\mathbf{Z}\longrightarrow\mathbf{W}$, such that $\mathbf{Z}_u\longrightarrow\mathbf{W}_u$ is the $\mathbf{E}_u$- minimal model of $\mathbf{W}_u$. Hence the E-minimal model of W and the E_0-minimal model of W_0 form a family. Now the process of constructing a divisorial contraction described in 3.3 can be done in families and therefore the divisorial contractions $Y\longrightarrow X$ and $Y_0\longrightarrow X_0$ form a family. Therefore there exists a family $\mathbf{Y}\longrightarrow\Delta$ such that $\mathbf{Y}_0\cong Y_0$, and $\mathbf{Y}_u\cong Y$ for all $u\neq 0$ in Δ. Hence if Y_0 is terminal, then by [**16**] Y is also terminal.

At this point I would like to mention that all the arguments so far work for the general D_{2n+1} case. However I do not know how to show that Y deforms to Y_0 for $n\geq 4$. I believe that a more carefull look at theorem 2.7 will treat the general case but the amount of calculations involved exceed the value of the result.

We now proceed to show that Y_0 is indeed terminal. The main point is to construct $h_0\colon Z_0\longrightarrow W_0$ explicitely and to show that

(i) No h_0-exceptional curve is contained in F^Z and therefore $F^Z\cong\mathbb{P}^2$. Then there are no flips involved and F^Z is contracted to a terminal singularity, obtaining Y_0.

(ii) Z_0 has isolated terminal singularities away from E_2^Z and hence W_0 is terminal.

We can construct Z_0 as follows. There is a sequence of crepant blow ups

$$W_4\xrightarrow{h_4}W_3\xrightarrow{h_3}W_2\xrightarrow{h_2}W_1\xrightarrow{h_1}W_0,$$

where h_i is the blow up of W_i along the birational transform of E in W_i. It is a straightforward calculation to check that W_4 has terminal hypersurface singularities. The point is that at the generic point of $E\cap F$, W_0 is an A_4 DuVal singularity. Then the process described above is just the blow up of a line through an A_4 DuVal singularity that intersects an edge of the dual graph. This leads to a crepant resolution of the singularity and hence W_4 has isolated terminal hypersurface singularities. Then in order to make E^{W_4} Cartier, all we need to do is blow it up. So let $W_4'\xrightarrow{h_4'}W_4$ be the blow up of E^{W_4}. Let $\psi=h_4'\circ h_4\circ h_3\circ h_2\circ h_1$. Then it is not difficult to check that no h_4'-exceptional curve is contained in $F^{W_4'}$. Contracting all the ψ-exceptional divisors we get the E-minimal model of W_0, Z_0 with all the properties claimed.

Case 2. None of the monomials yt and t^2 appear in $\phi_{\geq 2}(y,z,t)$. In this case we work explicitely with the equation of X and Γ. We use the normal form for X given in (7).

We start by describing $W = B_\Gamma X$. In the chart $x = xt$, $y = yt$ it is given by

$$x^2t + y^2tz + xz^2 + xtz\psi(z,t) + axt^{k+1} + \phi_{\geq 2}(yt,z,t) = 0.$$

Set $t = 0$ to see that E is given by $x = t = 0$, and F by $z = t = 0$. Let $L = E \cap F$: $x = z = t = 0$. Now it is easy to see that W is singular along L. This is what makes this case so different from the D_{2n} cases where W had only one singularity along L, and much more difficult to work.

As we said before, we try to show that none of the h_2-exceptional curves is contained in $F_1^{W_2}$ and also study the singularities of W_2 away from $F^{W_2} \cup F_1^{W_2}$. Moreover, Z can only have isolated cDV point over a cDV point and therefore we restrict our attention to what happens over non cDV points. One can also check the other charts and see that all the non cDV point are contained in the first one given by $x = xt$, $y = yt$. So it suffices to do all our calculations in that chart.

Let $h_1 : W_1 \longrightarrow W$ be the blow up of W along E. In the chart $x = xt$, W_1 is given by

$$x^2t^2 + y^2z + xz^2 + xzt\psi(z,t) + axt^{k+1} + \frac{1}{t}\phi_{\geq 2}(yt,z,t) = 0. \tag{3.2}$$

Let F_1 be the h_1-exceptional divisor. Setting $t = 0$ we see that

$$F_1 : z = t = 0 \tag{3.3}$$

$$E^{W_1} : t = y^2 + xz + \frac{1}{z}[\frac{1}{t}\phi_{\geq 2}(yt,z,t)]_{t=0} = 0. \tag{3.4}$$

Observe that F^{W_1} does not appear in this chart. Now write $\phi_2(y,z,t) = a_3zt$. Then $\frac{1}{t}\phi_2(yt,z,t) = a_3z$. Hence $[\frac{1}{t}\phi_{\geq 2}(yt,z,t)]_{t=0} = a_3z$. The singular locus of W_1 is contained in $F_1 \cap E^{W_1}$ which is the two lines, l_d, given by $z = t = y - d = 0$, with d such that $d^2 + a_3 = 0$. We will now study the singularities of W_1 along these lines. The change of variables $y \mapsto y + d$ brings the equations of l_d to $y = z = t = 0$, and

$$\begin{aligned} Y_1 : x^2t^2 + (y+d)^2z + xz^2 + xzt\psi(z,t) + axt^{k+1} \\ + \frac{1}{t}\phi_{\geq 2}((y+d)t,z,t) = 0. \end{aligned}$$

Look what happens along l_d by making the change of variables $x \mapsto x - \delta$. Y_1 is given by

$$\begin{aligned} (x-\delta)^2t^2 + (y+d)^2z + (x-\delta)z^2 + (x-\delta)zt\psi(z,t) \\ + a(x-\delta)t^{k+1} + a_3z + \frac{1}{t}\phi_{\geq 3}((y+d)t,z,t) = 0, \end{aligned}$$

and this gives

$$\begin{aligned} (x-\delta)^2t^2 + y^2z + 2dyz + (x-\delta)z^2 + (x-\delta)zt\psi(z,t) \\ + a(x-\delta)t^{k+1} + \frac{1}{t}\phi_{\geq 3}((y+d)t,z,t) = 0. \end{aligned} \tag{3.5}$$

It now follows that W_1 is singular along these two lines, and smooth away from them. Therefore we want to calculate W_2 in a neighborhood of these lines. Now it is possible to write

$$\frac{1}{t}\phi_{\geq 3}(yt,z,t) = z^2f(y,z) + t\phi(y,z,t)$$

and

$$\phi(y,z,t) = z\phi_1(y,z) + t\phi_2(y,z,t).$$

Then (8) becomes

$$\begin{aligned} x^2t^2+y^2z+xz^2+a_3z+xzt\psi(z,t)+axt^{k+1}& \\ +z^2f(y,z)+zt\phi_1(y,z)+t^2\phi_2(y,z,t)&=0. \end{aligned} \tag{3.6}$$

Moreover,

$$\begin{aligned} F_1 &: z=t=0 \\ E^{W_1} &: t=y^2+xz+a_3+zf(y,z)=0 \\ \cup l_d &: y^2+a_3=z=t=0 \end{aligned}$$

Now we want to describe W_2. It can be obtained by blowing up a suitable multiple of E^{W_1}. To find it look what happens along l_d. The change of variables $y \mapsto y+d$ and $x \mapsto x-\delta$ shows that the ideal defining $2E^{W_1}$

$$I_{2E^{W_1}} = (t^2, y^2+xz+a_3+xt\psi(z,t)+zf(y,z)+t\phi_1(y,z))$$

is principal at all but finitely many points of l_d. Therefore we can get W_2 by blowing up $2E^{W_1}$. At this point I want to say that the reason of the condition that yt does not exist in the equation of X is that if it appears then I don't see which multiple of E^{W_1} has to be blown up to get W_2. Now $W_2 \subset \mathbb{C}^4_{x,y,z,t} \times \mathbb{P}^1_{u,w}$ is given by the equations

$$wt^2-u[y^2+xz+zf(y,z)+xt\psi(z,t)+t\phi_1(y,z)]=0 \tag{3.7}$$

$$\begin{aligned} x^2t^2+y^2z+xz^2+a_3z+xzt\psi(z,t)+axt^{k+1}& \\ +z^2f(y,z)+zt\phi_1(y,z)+t^2\phi_2(y,z,t)&=0. \end{aligned} \tag{3.8}$$

In the affine chart $u=1$, W_2 is given by

$$y^2+xz+zf(y,z)+xt\psi(z,t)+t\phi_1(y,z)-wt^2=0 \tag{3.9}$$

$$\Phi(y,z,t)=x^2+wz+axt^{k-1}+\phi_2(y,z,t)=0. \tag{3.10}$$

Now we study what happens over l_d. The curves $C=h_2^{-1}(l_d)$ are given by

$$\begin{aligned} y^2+a_3=z=t&=0 \\ x^2+axt^{k-1}\mid_{t=0}+\phi_2(y,0,0)&=0. \end{aligned} \tag{3.11}$$

From the description of W_2 it follows that no component of C is contained in $F_1^{W_2}$. Next we want to see what kind of singularities W_2 has along C. The jacobian, J, of W_2 along C is

$$J = \begin{pmatrix} 0 & 2d & x_0+f(0,0) & x_0\psi(0,0)+\phi_1(0,0) & 0 \\ 2x_0+at^{k-1}|_{t=0} & \dfrac{\partial\phi_2}{\partial y}(d,0,0) & w+\dfrac{\partial\phi_2}{\partial z}(d,0,0) & \dfrac{\partial\Phi}{\partial t}(d,0,0) & 0 \end{pmatrix}$$

where $d^2+a_3=0$ and $x_0^2+at^{k-1}|_{t=0}x_0+\phi_2(d,0,0)=0$. If $d \neq 0$ then from (12) it follows that W_1 is cA_n along l_d and therefore W_2 can have at worst isolated cDV points on C. Hence the case of interest is when $d=0$ which implies that $a_3=0$ as well.

Now $x_0\psi(0,0)+\phi_1(0,0)$ is the coefficient of zt in (12) at any point on l given by $x \mapsto x+x_0$. Moreover, $x_0^2+at^{k-1}|_{t=0}x_0+\phi_2(0,0,0)$ is the coefficient of t^2. Hence if it is nonzero then W_1 is cA_n under C, and therefore W_2 has isolated cDV points along C. So we want to investigate what happens if $x_0\psi(0,0)+\phi_1(0,0)=0$. Now by looking at the Jacobian matrix J we see that W_2 is either singular along C or smooth. In the first case the resulting contraction will be only canonical and in the second we will have to see what happens in the other chart. W_2 is singular along C iff

$$\begin{gathered}
(2x_0 + at^{k-1}|_{t=0})(x_0 + f(0,0)) = 0 \qquad (3.12)\\
[(a(k-1)x_0t^{k-2})\ |_{t=0} + \frac{\partial \phi_2}{\partial t}(0,0,0)](x_0 + f(0,0)) = 0\\
\frac{\partial \phi_2}{\partial y}(0,0,0)(x_0 + f(0,0)) = 0\\
x_0\psi(0,0) + \phi_1(0,0) = 0\\
x_0^2 + ax_0t^{k-1}|_{t=0} + \phi_2(0,0,0) = 0
\end{gathered}$$

A similar calculation in the other affine piece of W_2 given by $w = 1$ leads to the same conclusion about the singularities of W_2 and we again obtain the equations (18).

Now we will show that there are no other h_2-exceptional curves that appear over the other affine piece of W_1 given by $t = tx$, and therefore the conditions given by (16) are necessary and sufficient. W_1 is given by

$$x^2t + y^2tz + z^2 + xzt\psi(z, xt) + ax^{k+1}t^{k+1} + \frac{1}{x}\phi_{\geq 2}(xyt, z, xt) = 0. \qquad (3.13)$$

Moreover, $x = 0$ gives that $F_1 : z = x = 0$, $F^{W_1} : z = t = 0$ and

$$E^{W_1} = (x, y^2t + z + \frac{1}{z}[\frac{1}{x}\phi_{\geq 2}(xyt, z, xt)]_{x=0}).$$

The singularities of W_1 lie on $L_1 \cup l_d = E^{W_1} \cap F_1$. The line $L_1 : x = z = t = 0$ is over L, and $l_d : x = y - d = t = 0$, with $d^2 + adt^{k-1}|_{t=0} + \phi_2(0,0,0) = 0$ as before, is over a point and we already studied the part away from zero in the other chart. So we only need to see what happens along L_1.
Claim: E^{W_1} is $\mathbb{Q}$-Cartier along L_1. Therefore there are no h_2-exceptional curves over L_1 and hence the conditions for nonexistence of a terminal contraction are precisely those given by (18).

We know that $2F^{W_1}$ is Cartier. Consider the index 1 cover of F^{W_1}

$$\pi : \tilde{W_1} \longrightarrow W_1.$$

We will show that $F_1^{\tilde{W_1}}$ is $\mathbb{Q}$-Cartier over L_1 and therefore F_1, and hence E^{W_1} too, is $\mathbb{Q}$-Cartier along L_1. By definition of the index 1 cover [**13**],

$$\tilde{W_1} = \mathrm{Spec}(\mathcal{O}_{W_2} \oplus I_{F^{W_1}, W_1})$$

It is easy to see now that $\tilde{W_1}$ is given by

$$x^2 + y^2uw + u^2 + xuw\psi(uw, xw^2) + bx^{k+1}w^{2k} + \frac{1}{xw^2}\phi_{\geq 2}(xyw^2, uw, xw^2) = 0 \qquad (3.14)$$

and π_1 is given by $t = w^2$ and $uw = z$. Moreover,

$$F_1^{\tilde{Y_1}} : u = x = 0$$

and therefore, since t^2 does not appear in $\phi_{\geq 2}(y,z,t)$, it is $\mathbb{Q}$-Cartier along $\pi_1^{-1}(L_1)$: $x = u = w = 0$. The claim now follows.

Therefore, a terminal contraction does not exist iff (18) are satisfied. Let $a_{i,j,k}$ be the coefficient of $y^i z^j t^k$ in $\phi_{\geq 2}(y,z,t)$. Then $f(0,0) = a_{0,2,1}$, $\frac{\partial \phi_2}{\partial t}(0,0,0) = a_{0,0,4}$, $\frac{\partial \phi_2}{\partial y}(0,0,0) = a_{1,0,2}$, $\phi_2(0,0,0) = a_{0,0,1}$ and $\phi_1(0,0) = a_{0,1,2}$. Substituting these to (18) we get the conditions claimed by the statement of the theorem.

We now want to give a geometric interpretetion to the existence of the quadratic part $\phi_2(y,z,t)$ in the equation of X.

Claim: $\phi_2(y,z,t) \neq 0$, iff the general hyperplane section of X through 0 is a D_4 DuVal singularity. In other words, $0 \in X$ is a cD_4 compound DuVal singularity.

Suppose that $\phi_2(y,z,t) \neq 0$. Then it is possible to deform X to an X_0 given by

$$x^2 + y^2 z + xz^2 + t\phi_2(y,z,t) = 0.$$

By assumption, the general hyperplane section S of X through Γ is a D_4 DuVal singularity. Therefore $0 \in X$ is cD_n for some n (If it was cA_n, then the general hyperplane section through Γ is A_m for some m [**20**]). The Milnor number is upper semicontinuous in families and hence if X_0 is cD_4, then so is X. Suppose that $t^2 \in \phi_2(y,z,t)$. Then deform X_0 to X_0' given by $x^2 + y^2 z + xz^2 + t^3 = 0$. Then the section $z = t$ is a D_4 DuVal singularity. We may do the same with the other monomial parts of $\phi_2(y,z,t)$ to see that indeed $0 \in X$ is cD_4.

Conversely, suppose that $0 \in X$ is cD_4. We will use the following property characterising D_4 DuVal singularities. Let $0 \in T$ be a D_n DuVal singularity. Then by using the Weierstrass preparation theorem, T is given by $x^2 + f(z,t) = 0$. This is D_4 iff the cubic part of the previous equation $f_3(z,t)$ is a product of three distinct linear forms.

So suppose that $\phi_2(y,z,t) = 0$ but $0 \in X$ is cD_4. Eliminating x in (7) we see that the cubic part of the equation of X is $y^2 z + t\phi_2(y,z,t)$. If $\phi_2(y,z,t) = 0$, then this can never be a product of three distinct linear forms.

Now suppose that Y is terminal. We want to find its index. Let S be the general hyperplane section of X contaning Γ and S' its birational transform in W. Then by [**20**] it follows that S' has exacty one singular point which is an A_4 DuVal singularity. So W is an A_4 DuVal singularity at the generic point of L and hence $5E$ and $5F$ are Cartier at the generic point of L. Arguing as in the proof of [**20**, Theorem 5.1] we conclude that

$$\text{index}(E^Z) = \text{index}(F^Z) = 5.$$

Note that the above arguments show that $Z = Z'$ and $F^Z \cong \mathbb{P}^2$. Let $a > 0$ such that

$$K_Z = h'^* K_Y + aF^Z. \tag{3.15}$$

Let $l \subset F^Z \cong \mathbb{P}^2$ be a general line. Then $l \cdot F^Z = -b/5$, with b a positive integer. We want to find b.

$$l \cdot (E^Z + 2F^Z) = l \cdot h^*(E + 2F) = -1. \tag{3.16}$$

Moreover, $l \cdot E^Z = 1/5(5E^Z) \cdot l$. This is easily computable. I claim that the scheme theoretic intersection of $5E^Z$ and F^Z is just $3L$. Then for general l, $l \cdot E^Z = 3/5$. To see the claim it suffices to work at the generic point of L. Then as we explained earlier, Z is an A_4 surface singularity there and E^Z, F^Z are two lines through the singular point. Moreover, by [**20**, Theorem 4.1, Proposition 4.6], it is possible to write at the generic point of L, Z as $xy - z^5 = 0$, E^Z is given by $x - z = y - z^4 = 0$, and F^Z by $x - z^2 = y - z^3 = 0$. A straightforward calculation shows that length$(5E^Z \cap F^Z) = 3$ (at the generic point of L), and hence $5E^Z \cap F^Z = 3L$. Therefore from (22) it follows that

$$l \cdot F^Z = 1/2(-1 - 3/5) = -4/5.$$

Now from (21) it follows that $a = 5/4$, and therefore Y has index 4.

Finally, the general section Σ of X through 0 is given by $z = ax + by + ct$. If one looks carefully through the calculations in the proof of theorem 3.6, it can be seen that the equations (18) are exactly the conditions for Σ not to be normal. Hence we get corollary 2.4. It is unfortunate that the only proof I know of this result is a computational one. □

Corollary 3.9. *Let $0 \in \Gamma \subset X$ as above. Let Σ be the general hyperplane section of X through 0. Write X as deformation of Σ, $X \longrightarrow \Delta$, using Γ as the parameter. There is a morphism $\Delta \xrightarrow{\phi} Def(\Sigma)$ inducing the above deformation. Then there is a subset Z of $Def(\Sigma)$, such that, a terminal divisorial contraction exists iff $Im(\phi) \subset Z$.*

Proof. Let $\Sigma' \xrightarrow{f} \Sigma$ be a birational morphism such that Σ' is normal, $-K_{\Sigma'}$ is f-ample, and Σ' admits a terminal 1-parameter deformation. Then there is a morphism $Def(\Sigma') \xrightarrow{\phi} Def(\Sigma)$ [**14**, Prop. 11.4]. Let $Z \subset Def(\Sigma)$ be the union of the images of all such morphisms. We may consider X as a deformation of Σ along Γ, say over a base Δ. Let $W \xrightarrow{g} X$ be the canonical contraction that always exists [**20**]. Then W can be considered as a deformation of $\Sigma' = g^*\Sigma$. If W is terminal, then we showed that Σ' is normal. Hence $Im[\Delta \longrightarrow Def(\Sigma)] \subset Z$. Conversely, if $Im[\Delta \longrightarrow Def(\Sigma)] \subset Z$, then there is a morphism $\Sigma' \longrightarrow \Sigma$ that lifts to a morphism $W \longrightarrow X$, where W is a terminal deformation of Σ', and is the required contraction. □

References

1. A. Corti, 'Factoring birational maps of threefolds after Sarkisov', *J. Algebraic Geometry* 4 (1995), 223-254.
2. A. Corti, M. Reid, 'Explicit birational geometry of 3-folds', *LMN* 281 (2000).
3. S. Cutkosky, 'Elementary contractions of Gorenstein threefolds', *Math. Ann.* 280 (1988), 521-525.
4. R. Hartshorne, 'Local Cohomology', *Lecture Notes in Mathematics* 41, Springer Verlag 1967.
5. M. Kawakita, 'Divisorial contractions in dimension 3 that contract divisors to compound cA_1 points', *Compositio Mathematica* 133 (2002), 95-116.
6. M. Kawakita, 'Divisorial contractions in dimension 3 that contract divisors to smooth points', *Invent. Math.* 145 (2001), 105-119.
7. Y. Kawamata, 'Crepant blowing up of 3-dimensional canonical singularities and its application to degenerations of surfaces', *Annals of Mathematics* 127 (1988), 93-163.
8. Y. Kawamata, 'Divisorial contractions to 3-dimensional terminal quotient singularities', in *Higher-dimensional complex varieties* (Trento 1994), 241-246, de Gruyter, Berlin, 1996.
9. Y. Kawamata, 'Deformations of canonical singularities', *J. Amer. Math. Soc.* 12 (1999), no 1, 85-92.
10. J. Kollár, 'Singularities of pairs', *Proc. Symp. Pure Math. Soc.* volume 62, 1997.
11. J. Kollár, 'Flips, flops, minimal models, etc.', *Surv. in Diff. Geom.* 1 (1992), 293-323.
12. J. Kollár, N. I. Shepherd-Barron, 'Threefolds and deformations of surface singularities', *Invent. Math.* 91 (1988), 299-338.

13. J. Kollár, S. Mori, ‘Birational geometry of algebraic varieties’, *Cambridge University Press*, 1988.

14. J. Kollár, S. Mori, ‘Classification of three-dimensional flips’, *J. Am.Math. Soc.* Vol. 5 no. 3 (1992), 533-703.

15. A. Markoe, H. Rossi, ‘Families of strongly pseudo convex manifolds’, *Symposium in several complex variables*, Lecture Notes in Math. vol 184, Springer, 1971, 182-208.

16. N. Nakayama, ‘Invariance of plurigenera of algebraic varieties’, *RIMS preprint* number 1191, 1998.

17. M. Reid, ‘Minimal models of canonical singularities’, *Advanced Studies in Pure Math.* 1, 1983, 131-180.

18. M. Reid, ‘Young person’s guide to canonical singularities’, *Proc. Symp. Pure Math. Soc.* volume 46, 1987.

19. V. Shokurov, ‘Prelimiting flips’, to appear in *Proc. of Steklov Institute*.

20. N. Tziolas, ‘Terminal 3-fold divisorial contractions of a surface to a curve I’, *Compositio Mathematica* 139, no 3, 239-261.

21. N. Tziolas, ‘Three dimensional divisorial extremal neighborhoods’, preprint.

22. N. Tziolas, ‘Families of D-minimal models and applications to 3-fold divisorial contractions’, *Proc. London Math. Soc.* (3) 90 (2005), 345-370.

Nikolaos Tziolas
Max Planck Institute für Mathematik
Vivatsgasse 7, Bonn, 53111,
Germany

tziolas@mpim-bonn.mpg.de

Current address:
Department of Mathematics
University of Crete
Knossos Ave, Heraklion 71409,
Greece

tziolas@math.uoc.gr